

Clique cycle-transversals in distance-hereditary graphs

Andreas Brandstädt¹,
 Simone Esposito²,
 Loana Tito Nogueira²,
 and
 Fábio Protti²

¹ Institut für Informatik, Universität Rostock, Germany,
 e-mail: ab@informatik.uni-rostock.de

² Instituto de Computação, Universidade Federal Fluminense, Niterói, RJ, Brazil,
 email: {loana,fabio}@ic.uff.br, simone.esposito@light.com.br

Abstract. A *cycle-transversal* of a graph G is a subset $T \subseteq V(G)$ such that $T \cap V(C) \neq \emptyset$ for every cycle C of G . A *clique cycle-transversal*, or *cct* for short, is a cycle-transversal which is a clique. Recognizing graphs which admit a cct can be done in polynomial time; however, no structural characterization of such graphs is known. We characterize distance-hereditary graphs admitting a cct in terms of forbidden induced subgraphs. This extends similar results for chordal graphs and cographs.

1 Introduction

A *cycle-transversal* of a graph G is a subset $T \subseteq V(G)$ such that $T \cap V(C) \neq \emptyset$ for every cycle C of G . When T is a clique, we say that T is a *clique cycle-transversal* or simply *cct*. A graph admits a cct if and only if it can be partitioned into a complete subgraph and a forest; by this reason such a graph is also called a $(\mathcal{C}, \mathcal{F})$ -graph in [3].

Finding a minimum cycle-transversal in a graph is NP-hard due to a general result in [12], which says that the problem of finding the minimum number of vertices of a graph G whose deletion results in a subgraph satisfying a hereditary property π on induced subgraphs is NP-hard. This result implies the NP-hardness of other problems involving cycle-transversals, for instance the problem of finding a minimum odd cycle-transversal (which is equivalent to finding a maximum induced bipartite subgraph), or the problem of finding a minimum triangle-transversal (which is equivalent to finding a maximum induced triangle-free subgraph). Odd cycle-transversals are interesting due to their connections to perfect graph theory; in [11], an $O(mn)$ algorithm is developed to find odd cycle-transversals with bounded size. In [8], the authors study the problem of finding C_k -transversals, for a fixed integer k , in graphs with bounded degree; among other results, they describe a polynomial-time algorithm for finding minimum C_4 -transversals in graphs with maximum degree three.

Graphs admitting a cct can be recognized in polynomial time, as follows. Note first that $(\mathcal{C}, \mathcal{F})$ -graphs form a subclass of $(2, 1)$ -graphs (graphs whose vertex set can be partitioned into two stable sets and one clique). The strategy for recognizing a $(\mathcal{C}, \mathcal{F})$ -graph G initially checks whether G is a $(2, 1)$ -graph, which can be done in polynomial time (see [2]). If so, then test, for each candidate clique Q of a $(2, 1)$ -partition of G , if $G - Q$ is acyclic (which can be done in linear time). If the test fails for all cliques Q , then G is not a $(\mathcal{C}, \mathcal{F})$ -graph, otherwise G is a $(\mathcal{C}, \mathcal{F})$ -graph. To conclude the argument, we claim that the number of candidate cliques Q is polynomial. Since G is a $(2, 1)$ -graph, let (B, Q) be a $(2, 1)$ -partition of $V(G)$ where B induces a bipartite subgraph and Q is a clique. Let (B', Q') be another $(2, 1)$ -partition of $V(G)$. Then $|Q' \setminus Q| \leq 2$ and $|Q \setminus Q'| \leq 2$, otherwise $G[B]$ or $G[B']$ would contain a triangle, which is impossible. Therefore, we can generate in polynomial time all the other candidate cliques Q' from Q . This is the same argument used to count sparse-dense partitions (for more details see [7]). Although recognizing graphs admitting a cct can be done in polynomial time, no structural characterization of such graphs is known, even for perfect graphs.

A similar sparse-dense partition argument can be employed to show that an interesting superclass of $(\mathcal{C}, \mathcal{F})$ -graphs, namely graphs admitting a clique triangle-transversal, can also be recognized in polynomial time. Such graphs are also known in the literature as $(1, 2)$ -split graphs. A characterization of this class is given in [13], where it has been proved that there are 350 minimal forbidden induced subgraphs for $(1, 2)$ -split graphs. When G is a perfect graph, being a $(1, 2)$ -split graph is equivalent to being a $(2, 1)$ -graph: observe that a perfect graph G contains a clique triangle-transversal if and only if G contains a clique that intersects all of its odd cycles. In [4] and [9], respectively, characterizations by forbidden induced subgraphs of cographs and chordal graphs which are $(1, 2)$ -split graphs are presented.

Deciding whether a distance-hereditary graph admits a cct can be done in linear-time using the clique-width approach, since the existence of a cct can be represented by a Monadic Second Order Logic (MSOL) formula using only predicates over vertex sets [6, 10]. However, no structural characterization for distance-hereditary graphs admitting a cct was known. In order to fill this gap, in this note we describe a characterization of distance-hereditary graphs with cct in terms of forbidden induced subgraphs.

2 Background

In this work, all graphs are finite, simple and undirected. Given a graph $G = (V(G), E(G))$, we denote by $\bar{G}$ the complement of G . For $V' \subseteq V(G)$, $G[V']$ denotes the subgraph of G induced by V' . Let $X = (V_X, E_X)$ and $Y = (V_Y, E_Y)$ be two graphs such that $V_X \cap V_Y = \emptyset$. The operations “+” and “ $\cup$ ” are defined as follows: the *disjoint union* $X \cup Y$, sometimes referred simply as *graph union*, is the graph with vertex set $V_X \cup V_Y$ and edge set $E_X \cup E_Y$; the *join* $X + Y$ is the graph with vertex set $V_X \cup V_Y$ and edge set

$$E_X \cup E_Y \cup \{xy \mid x \in V_X, y \in V_Y\}.$$

Let $N(x) = \{y \mid y \neq x \text{ and } xy \in E\}$ denote the open neighborhood of x and let $N[x] = \{x\} \cup N(x)$ denote the closed neighborhood of x . A *cut-vertex* is a vertex x such that $G[V \setminus \{x\}]$ has more connected components than G . A *block* (or *2-connected component*) of G is a maximal induced subgraph of G having no cut-vertex. A block is *nontrivial* if it contains a cycle; otherwise it is *trivial*.

For a set $\mathcal{F}$ of graphs, G is $\mathcal{F}$ -free if no induced subgraph of G is in $\mathcal{F}$.

Vertices x and y are *true twins* (*false twins*, respectively) in G if $N[x] = N[y]$ ($N(x) = N(y)$, respectively).

Adding a *true twin* (*false twin*, *pendant vertex*, respectively) y to vertex x in graph G means that for G and $y \notin V(G)$, a new graph G' is constructed with $V(G') = V(G) \cup \{y\}$ and $E(G') = E(G) \cup \{xy\} \cup \{uy \mid u \in N(x)\}$ ($E(G') = E(G) \cup \{uy \mid u \in N(x)\}$, $E(G') = E(G) \cup \{xy\}$, respectively).

The *complete* (resp. *edgeless*) graph with n vertices is denoted by K_n (respectively I_n). The graphs K_1 and K_3 are called *trivial graph* and *triangle*, respectively. The *chordless cycle* (*chordless path*, respectively) with n vertices is denoted by C_n (P_n , respectively). The graph C_n ($\overline{C_n}$, respectively) for $n \geq 5$ is a *hole* (*anti-hole*, respectively).

The *house* is the graph with vertices a, b, c, d, e and edges ab, bc, cd, ad, ae, be . The *gem* is the graph with vertices a, b, c, d, e and edges $ab, bc, cd, ae, be, ce, de$. The *domino* is the graph with vertices a, b, c, d, e, h and edges $ab, bc, cd, ad, be, eh, ch$.




Figure 1: House, hole, domino, and gem.

If H is an induced subgraph of G then we say that G *contains* H , otherwise G is *H-free*. A *clique* (resp. *stable* or *independent set*) is a subset of vertices inducing a complete (resp. edgeless) subgraph. A *universal vertex* is a vertex adjacent to all the other vertices of the graph. A *split graph* is a graph whose vertex set can be partitioned into a stable set and a clique. It is well known that G is a split graph if and only if G is $(2K_2, C_4, C_5)$ -free.

A *star* is a graph whose vertex set can be partitioned into a stable set and a universal vertex. A *bipartite graph* is a graph whose vertex set can be partitioned into two stable sets. A *cograph* is a graph containing no P_4 . A *chordal graph* is a graph containing no C_k , for $k \geq 4$. A *distance-hereditary graph* is a graph in which the distances in any connected induced subgraph are the same as they are in the original graph. A *threshold graph* is a graph that can be constructed from a one-vertex graph by repeated applications of the

following two operations: (a) addition of a single isolated vertex to the graph; (b) addition of a single universal vertex to the graph. It is well known that G is a threshold graph if and only if G is $(2K_2, C_4, P_4)$ -free.

If $T \cap V(C) \neq \emptyset$ for cycle C , we say that T *covers* C .

3 The forbidden subgraph characterization

The following well-known characterization of distance-hereditary graphs, also called *HHDG-free graphs*, will be fundamental for our result:

Theorem 1 [1] *The following are equivalent for any graph G :*

- (i) G is a distance-hereditary graph.
- (ii) G can be generated from a single vertex by repeatedly adding a pendant vertex, a false twin, or a true twin, respectively.
- (iii) G is (house, hole, domino, gem)-free. (See Figure 1.)

Let $G = (V, E)$ be a graph, and for a vertex $x \in V$, let $N^k(x) = \{y \in V \mid \text{dist}_G(x, y) = k\}$ for $k \geq 0$ denote the distance levels in G with respect to x . For $k = 2$, let $R = V \setminus (N[x] \cup N^2(x))$. The following are useful properties of distance-hereditary graphs:

Proposition 1 *Let G be distance hereditary and $u, v \in N^2(x)$.*

- (i) *If $uv \in E(G)$ or $uv \notin E(G)$ but connected by a path in $N^2(x) \cup R$ then $N(u) \cap N(x) = N(v) \cap N(x)$.*
- (ii) *If $N(x)$ is a stable set then for $u, v \in N^2(x)$ with $uv \notin E$, $N(u) \cap N(x)$ and $N(v) \cap N(x)$ do not overlap.*

Proof. (i): Since G is (house, hole)-free, u, v cannot have incomparable neighborhoods in $N(x)$. Moreover, since G is (house, hole, domino, gem)-free, the neighborhoods of u and v in $N(x)$ cannot properly contain one another, which shows Proposition 1.

(ii): Let us suppose, by contradiction, that u and v overlap. In this case, let a, b, c be vertices in $N(x)$ such that $au \in E(G)$, $av \notin E(G)$, $bu, bv \in E(G)$, $cu \notin E(G)$ and $cv \in E(G)$. Then $G[x, a, b, c, u, v]$ induces a domino, which is a contradiction. ■

Theorem 2 *Let G be a distance-hereditary graph. Then G admits a clique cycle transversal if and only if G is $(G_1, \dots, G_{12})$ -free.*




Figure 2: Forbidden subgraphs for distance-hereditary graphs with cct.

Proof. It is easy to see that $G_1, \dots, G_{12}$ from Figure 2 have no cct. For the converse direction, let G' be a distance-hereditary $(G_1, \dots, G_{12})$ -free graph. By Theorem 1, G' results, starting with a single vertex, by repeatedly applying one of the three operations in Theorem 1 (ii). Since adding a pendant vertex y to a vertex x in G does not create cycles with y , we can restrict ourselves to the following two cases: G' results from G by either adding a true twin or a false twin y to vertex x in G , and in both cases, we have to show that G' has a cct.

We can inductively assume that G has a cct Q . The vertex set $V(G)$ can be partitioned into $\{x\} \cup N(x) \cup N^2(x) \cup R$. Let $Q_1 = Q \cap N(x)$, $Q_2 = Q \cap N^2(x)$ and $N_1(x) = N(x) \setminus Q_1$.

3.1 Case 1: y is a true twin to x .

Let G' result from G by adding a true twin y to x in G . In this case, the possible cycles with y in G' are triangles xya for $a \in N(x)$, triangles yab for $a, b \in N(x)$, $ab \in E(G)$, and C_4 's $yabc$ for $a, b \in N(x)$, $ab \notin E(G)$, $c \in N^2(x)$. If $x \in Q$ or, more generally, $Q \subseteq N[x]$, then $Q \cup \{y\}$ is a cct of G' . Thus we have to consider the case $x \notin Q$. Since for a triangle yab also xab is a triangle which is covered by Q , the triangle yab is covered by Q , and similarly for the C_4 $yabc$ where $xabc$ is a C_4 in G covered by Q . Thus, we only have to

deal with triangles xya .

Claim 1 *If $x \notin Q$ then $G[N(x)]$ is a split graph with partition $(N_1(x), Q_1)$.*

Proof of Claim 1. For each edge $ab \in G[N(x)]$, xab is a triangle. Hence, a or b is in Q and $N_1(x)$ is a stable set. Since Q_1 is a clique, the claim follows. $\diamond$

Since G' is (G_1, G_2, G_3) -free, R induces a cycle-free subgraph in G .

Case 1.1: $G[N^2(x) \cup R]$ is cycle-free.

Claim 2 *If $x \notin Q$ then G' has a cct.*

Proof of Claim 2. Since Q is a clique and $G[N^2(x) \cup R]$ is cycle-free, Q_2 contains at most two vertices. If $Q_2 = \emptyset$ then $Q \cup \{y\}$ is a cct of G' . If a vertex $u \in Q_2$ has no neighbors in $N_1(x)$ then every cycle containing u also contains a vertex of Q_1 , i.e., $Q \setminus \{u\}$ is still a cct of G . Thus, assume without loss of generality that every vertex in Q_2 has a neighbor in $N_1(x)$.

If $Q_2 = \{u, v\}$, the neighborhood of Q_2 in $N_1(x)$ cannot contain two vertices a and b , otherwise by Claim 1 and Proposition 1 vertices x, y, a, b, u, v induce G_5 . Hence u and v have precisely one neighbor $a \in N_1(x)$ which must be adjacent to all vertices of Q_1 , otherwise if a misses a vertex $b \in Q_1$ then vertices x, y, a, b, u, v induce G_5 . Therefore $(Q \setminus \{u, v\}) \cup \{a, y\}$ is a cct of G' .

If $Q_2 = \{u\}$, we consider two subcases:

(i) Vertex u has a neighbor $a \in N_1(x)$ which misses some vertex $b \in Q_1$. Then every cycle C in G containing a, u but no vertex of Q_1 must also contain x . This is shown as follows: If C does not contain x then, by Proposition 1, C is either a triangle auv with $v \in N^2(x)$ or a C_4 $aucv$ with $c \in N_1(x), v \in N^2(x)$. In the former case by using Proposition 1, vertices x, y, a, b, u, v induce G_5 . The latter case cannot occur since the existence of cycle $aucv$ implies the existence of cycle $axcv$ in G , not covered by Q . This implies that $(Q \setminus \{u\}) \cup \{x, y\}$ is a cct of G' .

(ii) Every neighbor $a \in N_1(x)$ of u sees all vertices in Q_1 . Then $Q \cup \{a\}$ is a cct of G for some $a \in N_1(x)$ and, since by Claim 1 $N_1(x)$ is a stable set, every other neighbor $a' \in N_1(x)$ of u misses some vertex in $Q_1 \cup \{a\}$. By applying a similar argument as in (i), every cycle C in G containing a', u but no vertex of $Q_1 \cup \{a\}$ must also contain x . We conclude that $(Q \setminus \{u\}) \cup \{x, y, a\}$ is a cct of G' . This completes the proof of Claim 2. $\diamond$

Case 1.2: $G[N^2(x) \cup R]$ is not cycle-free.

We now assume that $G[N^2(x) \cup R]$ contains a cycle C . This cycle can be one of the following types (see Figure 3):

- (A₁) C has exactly one vertex u in $N^2(x)$, and C is a C_4 .
- (A₂) C has exactly one vertex u in $N^2(x)$, and C is a C_3 .
- (B₁) C has exactly two vertices u, v in $N^2(x)$, $uv \in E(G)$, and C is a C_4 .
- (B₂) C has exactly two vertices u, v in $N^2(x)$, $uv \in E(G)$, and C is a C_3 .
- (B₃) C has exactly two vertices u, v in $N^2(x)$, $uv \notin E(G)$, and C is a C_4 .
- (C₁) C is a C_4 with exactly three vertices u, v, w in $N^2(x)$ (which form a P_3 in $N^2(x)$).
- (D₁) C is a C_3 in $N^2(x)$.
- (D₂) C is a C_4 in $N^2(x)$.




Figure 3: Cycles in $G[N^2(x) \cup R]$.

Claim 3 $N(x)$ is a clique.

Proof of Claim 3. Suppose to the contrary that there are $a, b \in N(x)$ with $ab \notin E(G)$. Since G' is (G_1, G_2) -free, a and b must see each cycle in $G[N^2(x) \cup R]$. If C is a cycle of type (A₁) or (A₂) in $G[N^2(x) \cup R]$, i.e., with exactly one vertex u in $N^2(x)$ then a and b see u and we obtain G_7 or G_8 - contradiction. If C is of type (B₃) with $u, v \in N^2(x)$, $uv \notin E(G)$, then both a and b have to see C , and if not both a and b see both u and v then there is either a hole or domino or G_8 . Thus a and b see both u and v , i.e., there is G_{11} - contradiction.

We analyze the remaining cases by considering the following situation: If a sees vertex u and b sees vertex $v \neq u$ in $N^2(x)$ such that there exists a path linking u and v in $N^2(x)$ then by Proposition 1, a and b see a common edge $u'v'$; but then G' contains G_5 with x, y, a, b, u', v' - contradiction. This shows Claim 3. $\diamond$

Let $N_C^2(x)$ denote the set of all vertices in $N^2(x)$ which are contained in cycles of subgraph $G[N^2(x) \cup R]$. Since G is (G_1, G_2, G_3) -free, there is only one connected component in $G[N_C^2(x) \cup R]$. In addition, if $a \in N(x)$ then there is a triangle xya , and a must see every cycle in $G[N_C^2(x) \cup R]$.

Claim 4 *Every vertex in $N(x)$ sees every vertex in Q_2 .*

Proof of Claim 4. Since G' is (G_1, G_2, G_3) -free, any vertex $a \in N(x)$ sees at least one vertex u in each cycle of $G[N^2(x) \cup R]$. Since by Claim 3, $N(x)$ is a clique, and by Proposition 1, all vertices in Q_2 have the same neighborhood in $N(x)$, and vertex u sees a , all vertices in Q_2 see all vertices in $N(x)$ which shows Claim 4. $\diamond$

We conclude that if there is a cycle in $G[N^2(x) \cup R]$ and $x \notin Q$ then $N(x) \cup Q_2$ is a cct of G' , which finishes the proof in Case 1.

3.2 Case 2: y is a false twin to x .

Let G' result from G by adding a false twin y to x in G . We again inductively suppose that G has a cct Q . The possible cycles with y in G' are triangles yab for $a, b \in N(x)$, $ab \in E(G)$, C_4 's $yabc$ for $a, b \in N(x)$, $ab \notin E(G)$, $c \in N^2(x)$, and C_4 's $xyab$ for $a, b \in N(x)$.

If $|N(x)| = 1$ then Q is also a cct of G' . Now assume that $|N(x)| \geq 2$.

Recall that $V(G)$ is partitioned into $\{x\} \cup N(x) \cup N^2(x) \cup R$, and since G' is (G_1, G_2, G_3) -free, R induces a cycle-free subgraph in G' .

The fact below strengthens Claim 1.

Claim 5 $G'[N(x)]$ is a threshold graph.

Proof of Claim 5. Since G' is distance hereditary, $N(x)$ is P_4 -free, and since G' is G_5 - and G_6 -free, $N(x)$ is $2K_2$ - and C_4 -free, i.e., $G'[N(x)]$ is a threshold graph which shows Claim 5. $\diamond$

Case 2.1: $G[N^2(x) \cup R]$ is cycle-free.

We are going to show that also in this case, G' has a cct.

Recall that G has a cct Q , and let $Q_1 = Q \cap N(x)$, $Q_2 = Q \cap N^2(x)$, $N_1(x) = N(x) \setminus Q_1$. As in Claim 2, Q_2 can contain at most two vertices, and we can assume that every vertex in Q_2 has a neighbor in $N_1(x)$. Moreover, if $Q_2 \neq \emptyset$ then $x \notin Q$.

Case 2.1.1: $|Q_2| = 2$.

Let $Q_2 = \{u, v\}$; recall that by Proposition 1, u and v have the same neighborhood in $N_1(x)$. We distinguish between three subcases:

(i) If u, v have three neighbors a, b, c in (the stable set) $N_1(x)$ then vertices x, y, u, a, b, c (vertices x, y, v, a, b, c , respectively) induce G_4 in G' , which is impossible.

(ii) If u, v have exactly two neighbors a, b in $N_1(x)$ then there is no other vertex $c \in N_1(x)$, otherwise x, y, a, b, c, u, v induce G_{11} in G' . In addition, since G' is G_4 -free and by Claim 5, $G'[N(x)]$ is a threshold graph, either a or b sees all vertices of Q_1 , otherwise if a misses a' and b misses b' in Q_1 , respectively, then either G' contains G_4 (if we can choose $a' = b'$) or there is a P_4 $ab'a'b$ in $N(x)$. Suppose that a sees all vertices of Q_1 . Then every cycle in G' containing y also contains some vertex in $Q \cup \{a\}$, showing that $Q \cup \{a\}$ is a cct of G' .

(iii) If u, v have exactly one neighbor a in $N_1(x)$, we analyze the neighborhood of a . If a misses some vertex $b \in Q_1$ then there is no other vertex $c \in N_1(x)$ (otherwise x, y, a, b, c, u, v induce G_{11} if $bc \notin E(G)$ or c, b, x, u, a induce house if $bc \in E(G)$), and hence every cycle in G' containing y also contains some vertex in Q , i.e., Q is still a cct of G' . If a sees all vertices in Q_1 , every cycle containing u or v also contains some vertex in $Q_1 \cup \{a\}$, and hence $(Q \setminus \{u, v\}) \cup \{a, y\}$ is a cct of G' .

Case 2.1.2: $|Q_2| = 1$.

Let $Q_2 = \{u\}$. Since G' is G_4 -free, u has at most two neighbors in $N_1(x)$. If u has two neighbors a, b in $N_1(x)$ then, by Claim 5 and since G' is G_4 -free, one of them, say a , must see all vertices in Q_1 , and this means that every cycle containing u also contains some vertex in $Q_1 \cup \{a\}$ (recall that $G[N^2(x) \cup R]$ is cycle-free), i.e., $(Q \setminus \{u\}) \cup \{a, y\}$ is a cct of G' . If u has precisely one neighbor a in $N_1(x)$ and a sees all vertices in Q_1 , again $(Q \setminus \{u\}) \cup \{a, y\}$ is a cct of G' ; otherwise, a misses a vertex b in Q_1 , and the analysis is as follows:

(i') If $N_1(x)$ consists only of vertex a then every cycle in G' containing y also contains a vertex of Q_1 , and hence Q is a cct of G' .

(ii') If $N_1(x)$ contains a vertex $c \neq a$, we must have $bc \notin E(G)$ (otherwise x, a, b, c, u induce a house). We show that there is no cycle C in G containing a, u but no vertex of Q_1 . If there is such a cycle C then by Proposition 1, it must be a triangle auv with $v \in N^2(x)$, and then vertices x, y, a, b, c, u, v induce graph G_{11} , or it is a C_4 $auvw$ with $u, v \in N^2(x)$ and $w \in R$ but then there is a G_{12} or domino in G' . We conclude that $(Q \setminus \{u\}) \cup \{y\}$ is a cct of G' .

Case 2.1.3: $Q_2 = \emptyset$.

In this case, $Q \subseteq N[x]$. If there is a cct Q of G with $x \notin Q$ then $Q \cup \{y\}$ is a cct of G' and we are done. So we have to show that in Case 2.1.3, G has a cct Q with $Q \subseteq N(x)$.

In G , there are two types of cycles containing x : Triangles xab with $a, b \in N(x)$ and C_4 's $xabc$ with $a, b \in N(x)$ and $c \in N^2(x)$. Recall that by Claim 5, $N(x)$ induces a threshold graph and in particular is partitioned into a clique Q_1 and a stable set $N_1(x)$. Then Q_1 (and in general, every maximal clique in $N(x)$) covers every triangle xab since $ab \in E$.

The case of C_4 with x in G is more involved. Assume that there is a C_4 $xabc$ with

$a, b \in N(x)$ and $c \in N^2(x)$ which is not covered by Q_1 . If vertex a (vertex b , respectively) sees all vertices in Q_1 then the clique $Q_1 \cup \{a\}$ ($Q_1 \cup \{b\}$, respectively) covers abc as well. Otherwise both a and b have non-neighbors in Q_1 . Since by Claim 5, $N(x)$ is P_4 -free, a and b have a common non-neighbor, say d , in Q_1 . It follows that $cd \notin E$ (otherwise x, y, c, a, b, d induce G_4). If a and b miss another vertex $d' \in Q_1$ then x, y, d, d', a, b, c induce G_{11} , a contradiction. Thus one of a and b , say a , has at most one non-neighbor, say d , in Q_1 , and since $N(x)$ is a threshold graph, without loss of generality, the neighborhood of b in Q_1 is contained in the neighborhood of a in Q_1 ; in particular, b misses d , and, as above, c misses d . Let e be a neighbor of d in $N^2(x)$.

Claim 6 *Let e be a neighbor of d in $N^2(x)$. Then e misses a, b and c . (with a, b, c and d as described above).*

Proof of Claim 6.

We begin by observing that if $ce \in E$, by Proposition 1 (i), c and e have the same neighbors in $N(x)$ which is impossible since e sees d and c misses d . Therefore $ce \notin E(G)$. In this case, by Proposition 1 (ii), if e sees one of a and b , it must see both of them but now, x, y, e, a, b, d induce G_4 - a contradiction. Then e must also miss both a and b . $\diamond$

Suppose that $Q'_1 := (Q_1 \setminus \{d\}) \cup \{a\}$ is not a cct of G . Note that $N(x) \setminus Q'_1$ is stable. Then there is a cycle in G whose only vertex from Q_1 is d . Obviously, if C is a cycle containing d and an edge in $N(x)$ then Q'_1 covers C since $N(x) \setminus Q'_1$ is stable. Thus, we have to consider cycles without an edge in $N(x)$.

First consider a C_3 duv with $u, v \in N^2(x)$. Then by Claim 6, u and v miss a, b and c , and now, together with y , G' contains G_9 , a contradiction. If d is in a C_4 $duvw$ with $u, v \in N^2(x)$ and $w \in R$ then very similarly, together with y , G' contains G_{10} , a contradiction. Thus, d is not contained in any of such cycles.

If C is a C_4 with $d, z \in N(x)$ and $u, v \in N^2(x)$ then, again by Claim 6, u and v miss a, b and c . Then z must see a and b , otherwise there is a house or G_{12} in G' , together with y , but then $xadzu$ induce a house, a contradiction.

This also happens when d is in a C_4 with x and no vertex from Q'_1 . This final contradiction shows that in Case 2.1.3, there is a cct Q of G without x , and thus, there is a cct $Q \cup \{y\}$ in G' .

Case 2.2: $G[N^2(x) \cup R]$ is not cycle-free.

As in Case 1, we now assume that $G[N^2(x) \cup R]$ contains a cycle C (which implies that in this case, $x \notin Q$ holds).

Claim 7 $N(x)$ is I_3 -free.

Proof of Claim 7. Assume that $G[N^2(x) \cup R]$ is not cycle-free and $N(x)$ contains a stable set of three vertices a_1, a_2, a_3 . Let S be a maximal stable set in $N(x)$ containing a_1, a_2, a_3 .

Recall that by Proposition 1, for vertices $u, v \in N^2(x)$ in the same connected component of $G[N^2(x) \cup R]$, their neighborhoods in S are equal. In addition, no vertex $u \in N^2(x)$ sees at least three vertices in S , otherwise G_4 is contained in G' . Thus, for every pair of vertices $u, v \in N^2(x)$, $N(u) \cap S = N(v) \cap S \subseteq \{a_1, a_2\}$ holds.

If $u \in N^2(x)$ is in a cycle of type (A_1) or (A_2) then x, y, a_1, a_2, a_3, u and the vertices of the remaining cycle induce G_2, G_3, G_9 or G_{10} . If the cycle with $u, v \in N^2(x)$ is of type (B_1) , there is a house; if it is of type (B_2) or (D_1) , there is G_2 or G_{11} ; if of type (B_3) , there is G_3 or G_{12} ; and finally, if of type (C_1) or (D_2) , there is G_3 or G_{11} . This shows Claim 7. $\diamond$

We conclude that if y is a false twin to x and $G[N^2(x) \cup R]$ is not cycle-free then by Claim 7, $N_1(x)$ contains at most two vertices. If $|N_1(x)| \leq 1$ then Q is a cct of G' . If $N_1(x) = \{a, b\}$ then by Claims 5 and 7, one of them, say a , sees all vertices in Q_1 . By Proposition 1, either $Q_2 \cup \{a\}$ is a clique, and then $Q \cup \{a\}$ is a cct of G' , or a sees no vertex of Q_2 , and then let C be a cycle in $G[N^2(x) \cup R]$ and let $u \in V(C) \cap Q_2$. Since $xyab$ is a C_4 and G is (G_2, G_3) -free, there is an edge linking $xyab$ and C . By Proposition 1, we conclude that b sees all vertices in $C \cap N^2(x)$ and, therefore, in Q_2 . Now if there is some $a' \in Q_1$ then x, a, a', b, u induce either a house or a gem – a contradiction. Therefore $Q_1 = \emptyset$ and $Q'_2 \cup \{b\}$ is a cct of G' . This finishes the proof in Case 2 and thus also the proof of Theorem 2. $\blacksquare$

As a direct consequence of Theorem 2, we obtain another proof of a result in [3]:

Corollary 1 *If G is a cograph then G admits a clique cycle-transversal if and only if G is $(G_1, \dots, G_6)$ -free.*

Proof. Graphs G_1 to G_6 admit no cct. Conversely, G is also $(G_7, \dots, G_{12})$ -free (because all of them contain P_4). Since every cograph is a distance-hereditary graph, by Theorem 2 the corollary follows. $\blacksquare$

Corollary 2 *Let G be a distance-hereditary graph. Then G is a $(2, 1)$ -graph if and only if G is (G_1, G_5, G_6, G_7) -free.*

Proof. Graphs G_1, G_5, G_6 and G_7 are not $(2, 1)$ -graphs. Conversely, assume that G is (G_1, G_5, G_6, G_7) -free and G is not a $(2, 1)$ -graph. Let G' be a minimal induced subgraph of G which is not a $(2, 1)$ -graph. Note that being a $(2, 1)$ -graph is equivalent to admitting a clique that intersects every odd cycle. Thus G' does not admit a cct. By Theorem 2, G' is isomorphic to one of the graphs $G_1, G_2, \dots, G_{12}$. Since $G_2, G_3, G_4, G_8, G_9, G_{10}, G_{11}$, and G_{12} are $(2, 1)$ -graphs, it follows that G contains G_1, G_5, G_6 , or G_7 as an induced subgraph. $\blacksquare$

References

- [1] H.-J. Bandelt, H.M. Mulder. Distance-hereditary graphs. *Journal of Combinatorial Theory B* 41 (1986) 182-208.
- [2] A. Brandstädt. Partitions of graphs into one or two independent sets and cliques. *Discrete Mathematics* 152 (1996) 47-54. Corrigendum: *Discrete Mathematics* 186 (1998) 295.
- [3] A. Brandstädt, S. Brito, S. Klein, L. T. Nogueira, F. Protti. Cycle transversals in perfect graphs and cographs. Submitted manuscript, 2011.
- [4] R. S. F. Bravo, L. T. Nogueira, S. Klein. Cographs (k, l) -partitionable. *Proc. of the 7th International Colloquium on Graph Theory. Electronic Notes in Discrete Mathematics* 22 (2005) 277-280.
- [5] T. H. Cormen, C. E. Leiserson, R. L. Rivest, C. Stein. *Introduction to Algorithms*. MIT Press, 1990.
- [6] B. Courcelle, J. A. Makowsky, U. Rotics. Linear time solvable optimization problems on graphs of bounded clique-width. *Theory of Computing Systems* 33 (2000) 125–150.
- [7] T. Feder, P. Hell, S. Klein, R. Motwani. Complexity of Graph Partitions Problems. *31st Annual ACM Symposium on Theory of Computing (STOC 99)* (1999) 464-472.
- [8] M. Groshaus, P. Hell, S. Klein, L. T. Nogueira, F. Protti. Cycle Transversals in Bounded Degree Graphs. *Discrete Mathematics and Theoretical Computer Science* 13:1 (2011) 45–66.
- [9] P. Hell, S. Klein, L. T. Nogueira, F. Protti. Partitioning chordal graphs into independent sets and cliques. *Discrete Applied Mathematics* **141** (2004) 185–194.
- [10] M. Rao. MSOL partitioning problems on graphs of bounded treewidth and clique-width. *Theoretical Computer Science* 377 (2007) 260–267.
- [11] B. Reed, K. Smith and A. Vetta. Finding odd cycle transversals. *Operations Research Letters* 32 (2004) 299–301.
- [12] M. Yannakakis. Node- and edge-deletion NP-complete problems. *Proceedings of the Tenth Annual ACM Symposium on Theory of Computing – STOC’78*, pp. 253–264, 1978, ACM Press.
- [13] I. E. Zverovich, I. I. Zverovich. An Improvement of Gyárfás’ bounds on the maximal order of a minimal forbidden induced subgraph for $(1, 2)$ -split graphs. *Rutcor Research Report RRR* 37-2002.